

## An improved Altman type generalization of the Brézis–Browder ordering principle

ÁRPÁD SZÁZ\*

**Abstract.** *By using a modified argument, we prove an improvement of our former Altman type generalization of the Brézis–Browder ordering principle which yields a stronger maximum principle.*

**Key words:** *ordered sets, monotonicity and boundedness, maximal elements*

**AMS subject classifications:** Primary 06A06; Secondary 04A25

Received March 29, 2007

Accepted May 4, 2007

### Introduction

In 1976, to unify a number of diverse results in nonlinear functional analysis, H. Brézis and F. E. Browder [3] proved the following general ordering principle.

**Theorem 1.** *Let  $X$  be an ordered set; for  $x \in X$  denote  $S(x) = \{y \in X; y \geq x\}$ . Let  $\phi : X \rightarrow \mathbb{R}$  be a function satisfying*

- (1)  $x \leq y$  implies  $\phi(x) \leq \phi(y)$ ;
- (2) for any increasing sequence  $\{x_n\}$  in  $X$  such that  $\phi(x_n) \leq C < \infty$  for all  $n$ , there exists some  $y \in X$  such that  $x_n \leq y$  for all  $n$ ;
- (3) for every  $x \in X$  there exists  $u \in X$  such that  $x \leq u$  and  $\phi(x) < \phi(u)$ .

*Then, for each  $x \in X$ ,  $\phi(S(x))$  is unbounded.*

As a direct consequence of this theorem, the above authors derived the following maximum principle.

**Corollary 1.** *Let  $\phi : X \rightarrow \mathbb{R}$  be a function, bounded above, and satisfying*

- (1')  $x \leq y$  and  $x \neq y$  imply  $\phi(x) < \phi(y)$ ;
- (4) for any increasing sequence  $\{x_n\}$  in  $X$ , there exists some  $y \in X$  such that  $x_n \leq y$  for all  $n$ .

*Then, for each  $a \in X$ , there exists some  $\bar{a} \in X$  such that  $a \leq \bar{a}$  and  $\bar{a}$  is maximal (i. e.,  $S(\bar{a}) = \{\bar{a}\}$ ).*

---

\*Institute of Mathematics, University of Debrecen, H-4010 Debrecen, Pf. 12, Hungary, e-mail: [szaz@math.klte.hu](mailto:szaz@math.klte.hu)

The importance of this corollary lies mainly in the fact that it easily yields a simplified version of Ekeland's variational principle and hence also of Caristi's fixed point theorem. Moreover, it can also be used to prove Danes' drop theorem [3].

In 1982, having in mind the function  $\Phi$ , defined by  $\Phi(x, y) = \phi(x) - \phi(y)$  for all  $x, y \in X$ , M. Altman [1] generalized the above theorem in the following less satisfactory form.

**Theorem 2.** *Let  $(X, \leq)$  be an ordered set such that every totally ordered sequence  $\{x_n\} \subset X$  such that  $x_{n+1} \leq x_n$  for  $n = 1, 2, \dots$  has a minorant, i. e., there exists an element  $y \in X$  such that*

$$(i) \quad y \leq x_n \quad \text{for } n = 1, 2, \dots$$

*Let  $w = \Phi(x, y)$  be a real-valued function defined for all  $x, y \in X$  such that for each given  $y$ ,  $\Phi(\cdot, y)$  is bounded from below on  $S(y) = \{z \in X \mid z \leq y\}$ ;*

$$(ii) \quad \Phi(x, y) \leq 0 \quad \text{if } x \leq y \quad \text{for all } x, y \in X;$$

*(iii)  $\Phi$  is non-increasing in the second variable, i. e., for any given  $x \in X$ ,  $\Phi(x, y_2) \leq \Phi(x, y_1)$  if  $y_1 \leq y_2$  for all  $y_1, y_2 \in X$ ;*

$$(iv) \quad \liminf \Phi(x_{n+1}, x_n) = 0.$$

*Then for each  $x \in X$  there exists a  $y \in X$  such that  $y \leq x$  and  $z \leq y$  implies  $\Phi(z, y) = 0$ .*

As a direct consequence of this Theorem 2, the above author derived the following

**Corollary 2.** *Suppose that the hypotheses of Theorem 2 are satisfied with the assumption (ii) replaced by the stronger one*

$$(iib) \quad x \leq y \quad \text{and } x \neq y \quad \text{imply } \Phi(x, y) < 0.$$

*Then for each  $x \in X$  there exists  $\bar{x} \in X$  such that  $\bar{x} \leq x$  and  $\bar{x}$  is minimal, i. e.,  $z \leq \bar{x}$  implies  $z = \bar{x}$ .*

In 1984, M. Turinici [19] gave a better formulation and a metric generalization of the above theorem which also yields a maximum principle. Altman's theorem, in a somewhat improved form, has also been included in Zeidler [23, p. 515].

In 2001, not being aware of the works of M. Turinici, the present author also proved a generalization of Altman's theorem and derived a maximum principle. However, it has turned out that this theorem also contained several superfluous hypotheses.

Therefore, in the present paper we shall show that, by using a somewhat modified argument, we can actually prove a stronger result which may have a wider range of applications. For this, it is convenient to introduce some particular terminology.

## 1. Some general definitions

**Definition 1.** *If  $X$  is a set, then a function  $\Phi$  of  $X^2$  into  $\overline{\mathbb{R}}$  will be called an écart on  $X$ .*

**Example 1.** If  $\varphi$  and  $\psi$  are functions of  $X$  into  $\mathbb{R}$ , then the function  $\Phi$ , defined by  $\Phi(x, y) = \varphi(y) - \psi(x)$  for all  $x, y \in X$ , is a natural écart on  $X$ .

**Definition 2.** A set  $X$  equipped with a relation  $\leq$  will be called a goset (generalized ordered set).

**Remark 1.** A goset  $X$  will be called reflexive, symmetric and transitive, if the relation in it has the corresponding property.

**Definition 3.** If  $\Phi$  is an écart on a goset  $X$ , then the function  $\gamma_\Phi$ , defined by

$$\gamma_\Phi(x) = \sup_{y \geq x} \Phi(x, y)$$

for all  $x \in X$ , will be called the gauge of  $\Phi$ .

**Remark 2.** Note that if  $X$  is a reflexive goset and  $\Phi$  is as in Example 1, then  $-\infty < \gamma_\Phi(x)$  for all  $x \in X$ . Moreover, if  $a \in X$  is such that  $\varphi$  is bounded above on  $[a, +\infty[ = \{x \in X : a \leq x\}$ , then  $\gamma_\Phi(a) < +\infty$ .

Concerning the function  $\gamma_\Phi$ , it is also worth noticing the following

**Proposition 1.** If  $\Phi$  is an écart on a goset  $X$  such that for any  $x_1, x_2, y \in X$ , with  $x_1 \leq x_2$  and  $x_2 \leq y$ , there exists  $z \in X$ , with  $x_1 \leq z$ , such that  $\Phi(x_2, y) \leq \Phi(x_1, z)$ , then  $\gamma_\Phi$  is decreasing.

**Proof.** Suppose that  $x_1, x_2 \in X$  such that  $x_1 \leq x_2$ . If  $x_2 \not\leq y$  for all  $y \in X$ , then because of  $\sup(\emptyset) = -\infty$  we have  $\gamma_\Phi(x_2) = -\infty$ . Therefore,  $\gamma_\Phi(x_2) \leq \gamma_\Phi(x_1)$  automatically holds.

If  $y \in X$  such that  $x_2 \leq y$ , then by the assumption of the theorem there exists  $z \in X$ , with  $x_1 \leq z$  such that  $\Phi(x_2, y) \leq \Phi(x_1, z)$ . Hence, by the definition of the supremum, it is clear that

$$\Phi(x_2, y) \leq \Phi(x_1, z) \leq \sup_{w \geq x_1} \Phi(x_1, w) = \gamma_\Phi(x_1).$$

Therefore, by the definition of the supremum,  $\gamma_\Phi(x_2) = \sup_{y \geq x_2} \Phi(x_2, y) \leq \gamma_\Phi(x_1)$  also holds.  $\square$

Now, as an immediate consequence of the above proposition, we can also state

**Corollary 3.** If  $\Phi$  is an écart on a transitive goset  $X$  such that for any  $x_1, x_2, y \in X$ , with  $x_1 \leq x_2$  and  $x_2 \leq y$ , we have  $\Phi(x_2, y) \leq \Phi(x_1, y)$ , then  $\gamma_\Phi$  is decreasing.

**Remark 3.** Note that if  $X$  is a transitive goset and  $\Phi$  is as in Example 1 such that  $\psi$  is increasing, then  $\gamma_\Phi$  is already decreasing by the above corollary.

## 2. A generalized ordering principle

The importance of the above observations on  $\gamma_\Phi$  lies mainly in the following

**Lemma 1.** If  $\Phi$  is an écart on a goset  $X$  such that

- (1)  $\gamma_\Phi$  is decreasing;
- (2)  $-\infty < \gamma_\Phi(x)$  for all  $x \in X$ ;

(3)  $\gamma_{\Phi}(a) < +\infty$  for some  $a \in X$ ;

then there exists an increasing sequence  $(x_n)_{n=1}^{\infty}$  in  $X$ , with  $x_1 = a$ , such that

$$\lim_{n \rightarrow \infty} \gamma_{\Phi}(x_n) = \lim_{n \rightarrow \infty} \Phi(x_n, x_{n+1}).$$

**Proof.** Define  $x_1 = a$ . Then, by (2) and (3), we have  $-\infty < \gamma_{\Phi}(x_1) < +\infty$ . Therefore,

$$\gamma_{\Phi}(x_1) - 1 < \gamma_{\Phi}(x_1) = \sup_{y \geq x_1} \Phi(x_1, y).$$

Hence, by the definition of the supremum, it is clear that there exists  $x_2 \in X$ , with  $x_1 \leq x_2$ , such that

$$\gamma_{\Phi}(x_1) - 1 < \Phi(x_1, x_2).$$

Moreover, by using (2) and (1), we can also note that  $-\infty < \gamma_{\Phi}(x_2) \leq \gamma_{\Phi}(x_1) < +\infty$ . Therefore,

$$\gamma_{\Phi}(x_2) - 2^{-1} < \gamma_{\Phi}(x_2) = \sup_{y \geq x_2} \Phi(x_2, y).$$

Hence, by the definition of the supremum, it is clear that there exists  $x_3 \in X$ , with  $x_2 \leq x_3$ , such that

$$\gamma_{\Phi}(x_2) - 2^{-1} < \Phi(x_2, x_3).$$

Moreover, by using (2) and (1), we can note that  $-\infty < \gamma_{\Phi}(x_3) \leq \gamma_{\Phi}(x_2) < +\infty$ .

Now, by induction, it is clear that there exists an increasing sequence  $(x_n)_{n=1}^{\infty}$  in  $X$ , with  $x_1 = a$ , such that

$$\gamma_{\Phi}(x_n) - n^{-1} < \Phi(x_n, x_{n+1})$$

for all  $n \in \mathbb{N}$ . Moreover, we can also note that

$$\Phi(x_n, x_{n+1}) \leq \sup_{y \geq x_n} \Phi(x_n, y) = \gamma_{\Phi}(x_n)$$

for all  $n \in \mathbb{N}$ . Therefore, we actually have

$$\gamma_{\Phi}(x_n) - n^{-1} < \Phi(x_n, x_{n+1}) \leq \gamma_{\Phi}(x_n)$$

for all  $n \in \mathbb{N}$ . Hence, by using the monotonicity of the sequence  $(\gamma_{\Phi}(x_n))_{n=1}^{\infty}$  and some basic theorems on the limits of sequences in  $\overline{\mathbb{R}}$ , we can infer that

$$\lim_{n \rightarrow \infty} \gamma_{\Phi}(x_n) = \lim_{n \rightarrow \infty} \Phi(x_n, x_{n+1}).$$

□

Now, by using the above lemma, we can easily prove the following generalized ordering principle.

**Theorem 3.** *If  $\Phi$  is as in Lemma 1 and  $\alpha \in \overline{\mathbb{R}}$  such that*

*(4) each increasing sequence  $(x_n)_{n=1}^\infty$  in  $X$ , with  $x_1 = a$  is bounded above and satisfies*

$$\underline{\lim}_{n \rightarrow \infty} \Phi(x_n, x_{n+1}) \leq \alpha;$$

*then there exists  $b \in X$ , with  $a \leq b$ , such that  $\gamma_\Phi(b) \leq \alpha$ .*

**Proof.** If  $(x_n)_{n=1}^\infty$  is as Lemma 1, then by (4) we have

$$\lim_{n \rightarrow \infty} \gamma_\Phi(x_n) = \lim_{n \rightarrow \infty} \Phi(x_n, x_{n+1}) = \underline{\lim}_{n \rightarrow \infty} \Phi(x_n, x_{n+1}) \leq \alpha.$$

Moreover, by (4), there exists  $b \in X$  such that  $x_n \leq b$  for all  $n \in \mathbb{N}$ . Thus, in particular  $a = x_1 \leq b$ . Moreover, by (1) it is clear that  $\gamma_\Phi(b) \leq \gamma_\Phi(x_n)$  for all  $n \in \mathbb{N}$ , and thus

$$\gamma_\Phi(b) \leq \lim_{n \rightarrow \infty} \gamma_\Phi(x_n) \leq \alpha.$$

□

### 3. Applications of the generalized ordering principle

Theorem 3 easily yields the following extension of the main ordering principle of our former paper [13].

**Theorem 4.** *Assume that  $\Phi$  is an écart on a goset  $X$  such that  $\gamma_\Phi$  is decreasing. Moreover, assume that there exists  $\alpha \in \overline{\mathbb{R}}$  such that*

*(a)  $\alpha < \gamma_\Phi(x)$  for all  $x \in X$ ;*

*(b) each increasing sequence  $(x_n)_{n=1}^\infty$  in  $X$ , with  $\sup_{x_n \geq x_1} \Phi(x_1, x_n) < +\infty$ ,*

*is bounded above and satisfies  $\underline{\lim}_{n \rightarrow \infty} \Phi(x_n, x_{n+1}) \leq \alpha$ .*

*Then, we have  $\gamma_\Phi(x) = +\infty$  for all  $x \in X$ .*

**Proof.** If the required assertion is not true, then there exists  $a \in X$  such that  $\gamma_\Phi(a) < +\infty$ . Hence, it is clear that for any sequence  $(x_n)_{n=1}^\infty$  in  $X$ , with  $x_1 = a$ , we have

$$\sup_{x_n \geq x_1} \Phi(x_1, x_n) \leq \sup_{y \geq x_1} \Phi(x_1, y) = \gamma_\Phi(x_1) = \gamma_\Phi(a) < +\infty.$$

Therefore, by condition (b) and Theorem 3, there exists  $b \in X$  such that  $\gamma_\Phi(b) \leq \alpha$ . Moreover, by condition (a), we have  $\alpha < \gamma_\Phi(b)$ . This contradiction proves the required assertion. □

By using Theorem 3, we can also easily establish an extension of the main maximum principle of our former paper [13]. For this, it seems convenient to introduce the following

**Definition 4.** *An écart  $\Phi$  on a goset  $X$ , satisfying (1)–(3), will be called admissible at the point  $a$  if there exists  $\alpha \in \overline{\mathbb{R}}$  such that, in addition to (4), we also have*

(5)  $\alpha < \Phi(x, y)$  for all  $x, y \in X$  with  $x < y$ .

Now, by calling an element  $x$  of a goset  $X$  maximal if  $x \leq y$  implies  $x = y$  for all  $y \in X$ , we can easily state and prove the following generalized maximum principle.

**Theorem 5.** *If  $X$  is a goset and  $a \in X$  such that there exists an écart  $\Phi$  on  $X$  which is admissible at  $a$ , then there exists a maximal element  $b$  of  $X$  with  $a \leq b$ .*

**Proof.** By Definition 4, there exists  $\alpha \in \overline{\mathbb{R}}$  such that, in addition to (1)–(3), we also have (4) and (5). Thus, in particular by Theorem 3 there exists  $b \in X$ , with  $a \leq b$ , such that  $\gamma_{\Phi}(b) \leq \alpha$ , and thus  $\Phi(b, y) \leq \alpha$  for all  $y \in X$  with  $b \leq y$ .

Now, it remains only to show that  $b$  is maximal. For this, note that if this not the case, then there exists  $y \in X$ , with  $b \leq y$ , such that  $b \neq y$ , and thus  $b < y$ . Then, by the above property of  $b$ , we have  $\Phi(b, y) \leq \alpha$ . Moreover, by condition (5), we also have  $\alpha < \Phi(b, y)$ . This contradiction proves the maximality of  $b$ .  $\square$

**Remark 4.** *By making some obvious modifications in conditions (4) and (5), we can also easily establish the existence of an element  $b$  of  $X$ , with  $a \leq b$ , which is quasi-maximal in the sense that  $b \leq y$  implies  $y \leq b$  for all  $y \in X$ .*

*Note that if the goset  $X$  is reflexive, then every maximal element of  $X$  is quasi-maximal. While, if the goset  $X$  is antisymmetric, then the converse statement holds. Therefore, in a reflexive and antisymmetric goset the two notions coincide.*

**Acknowledgement.** The author is indebted to Zoltán Boros for some helpful and stimulating conversations which led to several improvements.

Moreover, the author also wishes to thank the referee for his valuable suggestions concerning formulations of remarks, shortenings of proofs, providing introduction and rewriting references. (Unfortunately, items [9], [11], [17], [18] and [21] were not available to the author.)

## References

- [1] M. ALTMAN, *A generalization of the Brézis-Browder principle on ordered sets*, Nonlinear Anal. **6**(1982), 157–165.
- [2] E. BISHOP, R. R. PHELPS, *The support functionals of a convex set*, In: *Convexity, Proc. Symp. Pure Math. VII* (V. Klee, Ed.), Amer. Math. Soc., 1963, 27–35.
- [3] H. BRÉZIS, F. E. BROWDER, *A general principle on ordered sets in nonlinear functional analysis*, Adv. Math. **21**(1976), 355–364.
- [4] A. BRØNDSTED, *On a lemma of Bishop and Phelps*, Pacific J. Math. **55**(1974), 335–341.
- [5] A. BRØNDSTED, *Common fixed points and partial order*, Proc. Amer. Math. Soc. **77**(1979), 365–368.

- [6] S. DANCs, M. HEGEDŰS, P. MEDVEGYEV, *A general ordering and fixed point principle in complete metric space*, Acta Sci. Math. (Szeged) **46**(1983), 381–388.
- [7] I. EKELAND, *On the variational principle*, J. Math. Anal. Appl. **47**(1974), 324–353.
- [8] I. EKELAND, *Nonconvex minimization problems*, Bull. Amer. Math. Soc. (N.S.) **1**(1979), 443–474.
- [9] A. HAMEL, *Variational Principles on Metric and Uniform Spaces* (Habilitation Thesis), Martin-Luther University, Halle-Wittenberg (Germany), 2005.
- [10] T. L. HICKS, *Some fixed point theorems*, Radovi Mat. **5**(1989), 115–119.
- [11] B. G. KANG, S. PARK, *On generalized ordering principles in nonlinear analysis*, Nonlinear Anal. **14**(1990), 159–165.
- [12] N. MIZOGUCHI, *A generalization of Brøndsted results and its applications*, Proc. Amer. Math. Soc. **108**(1990), 707–714.
- [13] Á. SZÁZ, *An Altman type generalization of the Brézis–Browder ordering principle*, Math. Moravica **5**(2001), 1–6.
- [14] Á. SZÁZ, *Lower and upper bounds in ordered sets without axioms*, Tech. Rep., Inst. Math., Univ. Debrecen **2004/1**, 1–11.
- [15] Á. SZÁZ, *Some easy to remember abstract forms of Ekeland’s variational principle and Caristi’s fixed point theorem*, Applicable Analysis and Discrete Mathematics **1**(2007), 335–339.
- [16] M. TURINICI, *Maximal elements in a class of order complete metric spaces*, Math. Japon. **25**(1980), 511–517.
- [17] M. TURINICI, *Maximality principles and mean value theorems*, An. Acad. Bras. Ciênc. **53**(1981), 653–655.
- [18] M. TURINICI, *A generalization of Brézis–Browder’s ordering principle*, A. Stint. Univ. Al. I. Cuza Iasi, Ser. Nouă, Mat. **28**(1982), 11–16.
- [19] M. TURINICI, *A generalization of Altman’s ordering principle*, Proc. Amer. Math. Soc. **90**(1984), 128–132.
- [20] M. TURINICI, *Metric variants of the Brézis–Browder ordering principle*, Demonstr. Math. **22**(1989), 213–228.
- [21] M. TURINICI, *Relational Brézis–Browder principles*, Fixed Point Theory **7**(2006), 111–126.
- [22] I. VÁLYI, *A general maximality principle and a fixed point theorem in uniform space*, Period. Math. Hung. **16**(1985), 127–134.
- [23] E. ZEIDLER, *Nonlinear Functional Analysis and its Applications I: Fixed-Point Theorems*, Springer-Verlag, New York, 1986.